

PRAZIMEC - D[®] Tablets for dogs (praziquantel, abamectin)

COMPOSITION

Content	Prazimec - D [®] Tablets for dogs
Praziquantel	0.050 g
Abamectin	0.002 g
Excipients	up to 0.480 g

PHARMACOLOGICAL ACTION

Praziquantel induces antihelminthic effect on cestodes (tapeworms) in dogs – *Echinococcus granulosus* (sexually mature), *Multiceps (Taenia) multiceps*, *T. hydatigena*, *T. ovis*, *T. pisiformis*, *T. taeniaeformis*, *Dipylidium caninum* (sexually mature and immature forms), *Diphyllobothrium latum*, as well as on some trematodes (flukes) as *Opisthorchis felineus*, *Paragonimus westermani*, *Apophallus donicum*, *Mesostephanus spp.* Abamectin is an endectocide which affects internal (nematodes) and external (arthropods – insects, mange mites and pasture ticks) parasites in dogs. Its antinematodal effect is directed against some roundworms – *Toxocara canis*, *Toxascaris leonina*, *Trichuris vulpis*, *Ancylostoma caninum*, *A. brasiliense*, *Uncinaria stenocephala*, *Strongyloides stercoralis*, and also against *Dirofilaria immitis*, *Filaroides osteri*. Abamectin induces insecticidal effect on some insects - *Linognathus setosus* (louse), *Ctenocephalides canis* (canine flea), *C. felis* (feline flea), *Pulex irritans* (human flea), as well as acaricidal effect on ticks – *Demodex canis*, *Otodectes cynotis*, *Sarcoptes canis*, *Notoedres spp.*, *Ixodes ricinus*, *Rhipicephalus sanguineus*, *Dermacentor reticulatus* and some other pasture ticks. The two active substances of the preparation complement each other's action, thereby providing a wider antiparasitic effect with a single administration. The product is well tolerated by dogs (with the exception of some breeds see under "Contraindications"). In case of overdosage, salivation, pupil dilation and gait disturbance may occur.

INDICATIONS

In infestation of dogs with endoparasites tapeworms – cestodes (teniae) and roundworms (nematodes); in filariasis of the heart; in infestations with fleas, lice, pasture ticks and mange mites including ear scabies; and particularly in cases of complex endo- and ectoparasitic invasion.

CONTRAINDICATIONS

Do not administer to dogs of Collie, Shelti and Bobtail breeds, to pregnant dogs, as well as to puppies below two months of age. Do not give more than four tablets as a single dose to dogs with a body weight exceeding 40 kg.

MODE OF ADMINISTRATION

Orally, with non-liquid feed.

TARGET SPECIES

Dogs.

DOSAGE

One tablet per 10 kg b. w.

The treatment may be repeated after a month. In cases of flea infestation the living premises must also be disinfected.

SIDE EFFECTS

Not established.

STORAGE

In the original packing, well closed, in dry and well-ventilated facilities, protected from direct sunlight.

PACKING

A box of 4 tablets.

WARNING

Keep out of the reach of children!

